

→ EARTH OBSERVATION SUMMER SCHOOL

Earth System Monitoring & Modelling

30 July–10 August 2018 | ESA–ESRIN | Frascati (Rome) Italy

Satellite Oceanography: an integrated perspective

ESA UNCLASSIFIED - For Official Use

How to contribute/accelerate new skills/discoveries to help reveal and model unknow unknowns from all available multi-modal satellite ocean remote sensing measurements ?

Some Earth Observation Challenges:

- *Upper vertical motions i.e. 3D dynamics (e.g. including time evolution) of the upper ocean,* Mesoscale and submesoscale circulation as key to control the vertical ocean pump and its impact on energy transport and biogeochemical cycles

- *Climate modelling due to these vast and diverse scales of fluid motions:* in the upper ocean, horizontal scales as big as basins and as small as cm-mm (capillary-gravity surface waves) contribute non-negligibly to air-sea exchanges and climate, and dynamics of scales of less than 30 km, is characterized by departures from the Earth's rotation constraint, i.e ageostrophic motions and strong impact of wind/wave transient forcings.

Problem: 1/100(0) year events now occur yearly!

Ocean remote sensing: a privileged view

- Spatially detailed
  - Spatial resolution from meters to Kms
  - A synoptic picture that is 100 km - 10 000 km wide
- Regularly repeated
  - Revisit intervals between 30 min. and 35 days
  - Continuously repeated over years to decades
- Global coverage
  - Satellites see the parts where ships rarely go
  - Single-sensor consistency - no intercalibration uncertainties
- Measures parameters that cannot be observed in situ
  - Surface roughness at short length scales (2-50 cm)
  - Surface slope (a few cm over 100s of kilometres)

New Era - Nanosatellites - CubeSat

A **CubeSat** is a type of miniaturized satellite for space research that usually has a volume of exactly 10 cm cube, and mass of no more than 1.33 kilograms.



ESA UNCLASSIFIED - For Official Use

Big Data & AI/Data Science

Data



Big data infrastructure

DL/AI

Deep Learning





In situ data

Remote sensing data



Simulation data

Numerous questions and challenges

Some of the Living Planet Challenges to better assess the existing pressures on the marine environment (e.g. overfishing, pollution, habitat destruction, ...) potentially leading to increased risks to global food security, economic prosperity, …

Evolution of coastal ocean systems including the interactions with land in response to natural and human-induced environmental perturbations

Mesoscale and submesoscale circulation and the role of the vertical ocean pump and its impact on energy transport and biogeochemical cycles

Response of the marine ecosystem and associated ecosystem services to natural and anthropogenic changes,

Physical and biogeochemical air/sea interaction processes on different spatio-temporal scales and their fundamental role in weather and climate

Sea level changes from global to coastal scales and from days (e.g. storm surges) to centuries (e.g. climate change)

Numerous questions and challenges

How can we map the distribution of marine plastic Debris?

Has the Agulhas current strengthened in the last 5 years?

Is the surface circulation of the Black Sea and in the Mediterranean Sea stable?

How is the Arctic Ocean changing ?

How is marine biodiversity changing, locally, regionally, globally ?

What is the extent of ocean acidification ?

Are western boundary currents changing, the Gulf Stream ?

How can ship routing be optimised ?

Why and where is regional sea level changing?

How are our coastal regions changing?

How can we map estuary systems from space?

- … most observations are not yet sufficiently explored and used

Synergy between high and medium resolution observations to reveal mean states and trends, near-surface ocean-atmosphere dynamics, local and non-local interactions, convergence/divergence surface fronts and numerous roughness contrasts

Atmospheric and Oceanic observations generally produce high quality data, but it is often too sparse (many gaps where information is missing, and/or often too local in both space and time)

How can we use observed data in combination with the physical knowledge of stochastic processes in nonlinear dynamical systems to estimate and model those effects on the variability of computationally resolvable scales of motion that are caused by the small, rapid, unresolvable scales of fluid motion that upscaling in data assimilation leaves out?

Climate models are too coarse to resolve clouds



Global model:
~100 km resolution

NASA MODIS

Cloud scales: ~10-100 m

SST

1 jour



MSG/SEVIRI (10km, 3 heures)



SAF O&SI NAR pour AVHRR17 (2km, 2 passes/jour)



SAF O&SI NAR18 pour AVHRR17 (2km, 2 passes/jour)



ATS



Avhrr 18



Avhrr 17



ENVISAT/AATSR (1 km, 14-15 orbites/jour)

Multi-satellite product

Sea surface temperature



Global SST


60°N
30°N
0°
30°S
60°S
150°W
120°W
90°W
60°W
30°W
0°
30°E
60°E
90°E
120°E
150°E
30
28
26
24
22
20
18
16
14
12
10
8
6
4
2
0
−2

Small scales / mesoscale

SST - Modis(L2P)



SST - AMSRE(L3)



Characterizing the submesoscale - Spectral approach





Schematic illustrating the different remote-sensing methods and classes of sensors used in satellite oceanography, along with their applications (from Robinson, 2004).





Main message (to be repeated) …

- Today ideal instrument … (wide-swath, high-resolution, topography, roughness, Doppler, emissivity, reflectance, …) = the combined use of observations, including in situ measurements
- Very (too) large number of spatio-temporal scales under local and non-local interactions
- Improved technologies (instruments, resolution, computer capabilities, storage, dissemination) all contribute to improved combined analysis
- Theoretical frameworks and numerical simulations can be used to assess the causes and contexts of the different observations (including sensor physics, observability conditions and instrument capabilities), to refine dynamical/statistical gap filling methods
- New challenges, new altimeter instruments (SARAL, Sentinel-3, SWOT, …, CubeSat opportunities), possible new high-resolution microwave instruments (10-20 km), Doppler measurements to infer sea surface currents (SAR and/or RAR-SKIM or DopplerScat), and combined roughness contrasts as local quantitative proxies to trace strong surface gradient areas

Satellite instruments generally measure 2D surface expressions of 4D structures



Typical Atmosphere Space-Time scales



Satellite synthetic aperture radar (SAR)

light = more short waves
~ stronger wind forcing
~ air-sea heat, KE, momentum, gas exchange

dark = weak wind or surfactant



February 19, 2015

Image: MPC Sentine-1 portal



strong cold wind over warmer ocean


strong
mean wind
ABL rolls
warm
cold
~1 km
~1 km
mixed layer
thermocline
SAR signature



proper atmosphere-wave-ocean coupling

We can illustrate this partitioning using the expression for the total momentum flux at the



Figure 2. Some of the processes that govern the transfer of heat, mass, and momentum within the coupled boundary layers.

ocean surface (i.e., where the turbulent component becomes negligible) derived by Deardorff (1967). Deardorff (1967) derived this expression by evaluating the integrated horizontal momentum equation at the ocean surface to obtain

$$\tau_a = \rho_a \left[-\overline{\tilde{u}\tilde{w}} + \nu \frac{\partial U}{\partial z} \right]_\eta \approx \overline{p_\eta \frac{\partial \eta}{\partial x}} + \rho_a \nu \frac{\partial U}{\partial z} \Big|_\eta = \tau_{aw} + \tau_{ao} \qquad \textbf{(15)}$$

where η is the wave height, p_η is the surface pressure, and the small component of the viscous stress associated with inclinations of the interface has been neglected. The stresses given on the RHS on based on the nomenclature given in Lionello et al. (1996, 1998), where τ_{aw} represents the momentum transfer from the wind to the waves (i.e., the wave-induced flux), while τ_{ao} represents the direct momentum transfer from the wind to the ocean (i.e., **viscous stress**). This

Schematic illustration of the Langmuir circulation (first described by Langmuir, 1938). The separation scale of the convergence zones are typically 10-100 m



Typical Ocean Space-Time Scales




Synoptic Wind
Swell Rays
Density Front
Inertial Motion
Stokes Drift
Internal Waves
Surface Roughness Anomaly
SQG
Wind Stress
Wind Waves
Surface
Mixing Layer Depth
1 day
10-30 days
1 year
Anti-cyclonic Eddy
Ekman Spiral
Ekman Transport
Turbulence
Langmuir Circulation
O(10m)
Internal Waves
Mixed Layer Depth
O(100m)
Maximum N
Remnant Layer Depth
O(1000m)
Density Profile
Summer
Heating
Winter
Depth


Ocean-Atmosphere
Interactions



Figure 1. Future role of wave models as an essential coupling component for ocean-atmosphere-carbon-cycle modets developed in the context ot the World Climate and Global Change programs.


Vertical scales
1000m
100m
10m
Dissipation
aux frontières
cascade KE
Circulation
Thermohaline
(100-1000ans)
Gyres
(10ans)
Mésoéchelle
(i.e. jets,
tourbillons d'outflow,...)
(qqs mois)
ondes de gravité internes
(qqs min-24h)
Sous-mésoéchelle
(i.e. filaments, fronts,...)
(qqs jours)
ondes inertielles
(24h)
cascade directe KE
turbulence stratifiée 3D
stratification domine
?
rotation domine
turbulence 3D
Dissipation
intérieure
1m
1km
100km
10⁴km
Horizontal scales

Satellite synthetic aperture radar (SAR)

light = more short waves
~ stronger wind forcing
~ air-sea heat, KE, momentum, gas exchange

dark = weak wind or surfactant



February 19, 2015

Image: MPC Sentine-1 portal

Inter-scale interactions

- dispersion of biochemical substances?
- High-resolution = Large Eddy Simulation (LES)



February 20, 2015

Image: MPC Sentine-1 portal


©ESA 2009
0
6
12 km



ESA UNCLASSIFIED - For Official Use



Landsat 8 false color image, August 11, 2015.
http://landsat.visibleearth.nasa.gov/view.php?id=86449.

Stirring and mixing : interplay and scale





Application to Oil Spills Detection

Small scales / mesoscale



Lagrangian advection to dynamically interpolate large-scale tracer (sea surface temperature field, left) onto a high-resolution product (right). Particle trajectories computed using altimetry-derived velocities (AVISO, weekly 1/3°) with 3 hours time steps



Small scales / mesoscale



Small scales / mesoscale



The blended satellite products allow to estimate the impact of surface currents on the biogeochemical transport, on the dispersion of pollutants and oil spills



Forecast of oil spill dispersion in the Gulf of Mexico on 25 june 2010: red and blue show regions of strong oil dispersion within 3 days. This diagnosis, based on altimetric data, compared well with what was observed (Mezic et al, Science, 2010).

However these satellite datasets (altimetric and microwave data) cannot capture ocean dynamics at scales smaller than 100 km because of the resolution (or/and noise level).

Observed data in combination with the physical knowledge of stochastic processes in nonlinear dynamical systems



LU :Adding random velocity

Large scales:

$\boldsymbol{w}$

Small scales:

$\boldsymbol{\sigma}\dot{\boldsymbol{B}}$

Variance tensor:

$\boldsymbol{a} = \boldsymbol{a}(\boldsymbol{x}, \boldsymbol{x}) = \frac{\mathbb{E}\{\boldsymbol{\sigma} d\boldsymbol{B}\,(\boldsymbol{\sigma} d\boldsymbol{B})^T\}}{dt}$

Resolved large scales

White-in-time small scales

$$\boldsymbol{v} = \boldsymbol{w} + \boldsymbol{\sigma}\dot{\boldsymbol{B}}$$

References :

Mikulevicius and Rozovskii, 2004
Flandoli, 2011

Memin, 2014
Resseguier et al. 2017 a, b, c
Chapron et al. 2017
Cai et al. 2017

Holm, 2015
Holm and Tyranowski, 2016
Arnaudon et al., 2017

Cotter and al 2017
Crisan et al., 2017
Gay-Balmaz & Holm 2017
Cotter and al 2018 a, b

Large scales:

$\boldsymbol{w}$

Small scales:

$\boldsymbol{\sigma}\dot{\boldsymbol{B}}$

Variance tensor:

$\boldsymbol{a} = \boldsymbol{a}(\boldsymbol{x}, \boldsymbol{x}) = \frac{\mathbb{E}\{\boldsymbol{\sigma} d\boldsymbol{B}\,(\boldsymbol{\sigma} d\boldsymbol{B})^T\}}{dt}$

Advection of tracer Θ

Multiplicative random forcing

Balanced energy exchanges

Advection

Diffusion

$$\partial_t \Theta + \boldsymbol{w}^\star \cdot \nabla\Theta + \boldsymbol{\sigma}\dot{\boldsymbol{B}} \cdot \nabla\Theta = \nabla \cdot \left(\frac{1}{2}\boldsymbol{a}\nabla\Theta\right)$$

Drift correction

Goal is a hierarchical system that integrates data and models (and can also be used to design observing systems)



Upper ocean atmospheric forcing monitoring



Upper ocean geostrophy monitoring



Upper ocean circulation monitoring



Upper ocean ageostrophic current monitoring



Total ocean surface current monitoring



- … most observations are not yet sufficiently explored and used

Synergy between high and medium resolution observations to reveal mean states and trends, near-surface ocean-atmosphere dynamics, local and non-local interactions, convergence/divergence surface fronts and numerous roughness contrasts

Atmospheric and Oceanic observations generally produce high quality data, but it is often too sparse (many gaps where information is missing, and/or often too local in both space and time)

How can we use observed data in combination with the physical knowledge of stochastic processes in nonlinear dynamical systems to estimate and model those effects on the variability of computationally resolvable scales of motion that are caused by the small, rapid, unresolvable scales of fluid motion that upscaling in data assimilation leaves out?